

Mechanism of Cellular Respiration:

Aerobic respiration is the process most living things undergo to use food energy. Here we will investigate the definition, the steps of the process, what goes in and what comes out of the process, and the chemical formula. Get ready to breathe!

Aerobic Respiration:

Take a deep breath...now exhale. Do you feel it? Every time you breathe in oxygen and exhale carbon dioxide, you are exchanging gases that are a crucial part of your energy metabolism. We don't normally equate oxygen and breathing with metabolizing food, yet the latter cannot occur without the former.

Most living organisms undergo this process, from single-celled bacteria to the multi-celled blue whale. While many microorganisms can accomplish this task within their single cell, we larger macroorganisms have evolved body organs dedicated to efficiently obtaining enough oxygen, and eliminating enough carbon dioxide, to undergo enough **aerobic respiration** to fuel our large, complex selves.


Definition of Aerobic Respiration:

Respiration is the metabolic process of most living things in which food molecules or glucose are turned into usable energy for the cell, called ATP.

Respiration is the anti-process to **photosynthesis**, the process in which plants use sunlight and carbon dioxide to build food molecules releasing oxygen as a waste product.

During **aerobic respiration**, oxygen is present and helps the process to crank out energy very efficiently. Some organisms can also undergo **anaerobic respiration**, in which oxygen is absent, and a somewhat less efficient method of metabolism takes place. While photosynthesis takes place in the chloroplasts of plant and algae

cells, aerobic respiration takes place in the **cytoplasm**, or the gooey inner cell space and mitochondria of all eukaryotic cells.




The Steps of Aerobic Respiration:

It all starts with a sugar! An organism takes in carbohydrates for energy, and the digestion process breaks the carbs down into their smallest units, **glucose**, a type of sugar molecule. Cells then make energy by breaking the glucose molecule down and releasing its electrons, which are later used to help crank out ATP. There are three main steps in this process.

There are three main steps in this process. It begins with **glycolysis**. During glycolysis, the 6-carbon glucose molecule undergoes a series of reactions that break it down into two 3-carbon **pyruvate molecules**. The purpose of this process is **to release electrons from the bonds in the glucose**, which are scooped up by an acceptor molecule called NAD^+ , turning it into NADH when it accepts the electrons. In this process, two molecules of ATP are made. This step occurs in the


cytoplasm, and the pyruvate and NADH molecules then enter the mitochondria for the next step.



The next phase of aerobic respiration is the **citric acid cycle**, also known as the **Kreb's cycle**, named for the biochemist who discovered it. To prepare for this stage, the pyruvate molecules from glycolysis are converted to a 2-carbon compound called **Acetyl CoA**. What happened to the third carbon? You just exhaled it in the form of carbon dioxide!

With each turn of the cycle, the Acetyl CoA is broken down and rebuilt into carbon chains. The purpose is to extract electrons from them and generate more ATP, similar to the more simple process of glycolysis. NAD^+ is used again to pick up the electrons released, as is another acceptor molecule, $FADH$, which becomes

FADH₂ when reduced. These acceptor molecules get loaded up with electrons, like cargo trucks, and carbon dioxide is released as the carbon chains are broken down and new Acetyl CoA comes in. Exhale!



Where do the cargo trucks go once they are loaded up? The FADH₂ and NADH molecules take their electrons to the inner mitochondrial membrane for the final stage of aerobic respiration, **oxidative phosphorylation**. Here, using the power of a concentration gradient, a very large amount of ATP is generated. The electrons from the citric acid cycle are dropped off and used to force hydrogen atoms that were released when the acceptor molecules picked up electrons in Steps 1 and 2, to pump against their concentration gradient. Proteins embedded in the membrane undergo active transport to push all these hydrogens into a highly concentrated

area, just so they can then rush downward through an enzyme, the **ATP Synthase**, which turns like a gear to crank out about 32 ATP!

To unlock this lesson you must be a Study.com Member. [Create your account](#)

A set of metabolic reactions and processes that take place in the cells of organisms to **convert biochemical energy from nutrients into adenosine triphosphate (ATP)**, and then release waste products.^[1] The reactions involved in respiration are catabolic reactions, which break large molecules into smaller ones, releasing energy in the process, as weak so-called "high-energy" bonds are replaced by stronger bonds in the products. Respiration is one of the key ways a cell releases chemical energy to fuel cellular activity. **Cellular respiration is considered an exothermic redox reaction which releases heat.** The overall reaction occurs in a series of biochemical steps, most of which are redox reactions themselves. Although technically, cellular respiration is a combustion reaction, it clearly does not resemble one when it occurs in a living cell because of the slow release of energy from the series of reactions.

Nutrients that are commonly used by animal and plant cells in respiration include **sugar, amino acids and fatty acids**, and the most common oxidizing agent (electron acceptor) is molecular oxygen (O₂). The chemical energy stored in ATP (its third phosphate group is weakly bonded to the rest of the molecule and is cheaply broken allowing stronger bonds to form, thereby transferring energy for use by the cell) can then be used to drive processes requiring energy, **including biosynthesis, locomotion or transportation of molecules across cell membranes.**

Aerobic respiration

Aerobic respiration (red arrows) is the main means by which both fungi and animals utilize chemical energy in the form of organic compounds that were previously created through photosynthesis (green arrow).

Aerobic respiration requires oxygen (O₂) in order to create ATP. Although carbohydrates, fats, and proteins are consumed as reactants, it is the preferred method of pyruvate breakdown in glycolysis and requires that pyruvate enter the mitochondria in order to be fully oxidized by the Krebs cycle. The products of this process are carbon dioxide and water, but the energy transferred is used to break bonds in ADP as the third phosphate group is added to form ATP (adenosine triphosphate), by substrate-level phosphorylation, NADH and FADH₂


The negative ΔG indicates that the reaction can occur spontaneously.

The potential of NADH and FADH₂ is converted to more ATP through an electron transport chain with oxygen as the "terminal electron acceptor". Most of the ATP produced by aerobic cellular respiration is made by oxidative phosphorylation. This works by the energy released in the consumption of pyruvate being used to create a chemiosmotic potential by pumping protons across a membrane. This potential is then used to drive ATP synthase and produce ATP from ADP and a phosphate group. Biology textbooks often state that 38 ATP molecules can be made per oxidised glucose molecule during cellular respiration (2 from glycolysis, 2 from the Krebs cycle, and about 34 from the electron transport system).^[2] However, this maximum yield is never

quite reached because of losses due to leaky membranes as well as the cost of moving pyruvate and ADP into the mitochondrial matrix, and current estimates range around 29 to 30 ATP per glucose.^[2]


Aerobic metabolism is up to 15 times more efficient than anaerobic metabolism (which yields 2 molecules ATP per 1 molecule glucose). However some anaerobic organisms, such as methanogens are able to continue with anaerobic respiration, yielding more ATP by using other inorganic molecules (not oxygen) as final electron acceptors in the electron transport chain. They share the initial pathway of glycolysis but aerobic metabolism continues with the Krebs cycle and oxidative phosphorylation. The post-glycolytic reactions take place in the mitochondria in eukaryotic cells, and in the cytoplasm in prokaryotic cells.

Glycolysis



Out of the cytoplasm it goes into the Krebs cycle with the acetyl CoA. It then mixes with CO₂ and makes 2 ATP, NADH, and FADH. From there the NADH and FADH go into the NADH reductase, which produces the enzyme. The NADH pulls the enzyme's electrons to send through the electron transport chain. The electron transport chain pulls H⁺ ions through the chain. From the electron transport chain, the released hydrogen ions make ADP for an end result of 32 ATP. O₂ attracts itself to the left over electron to make water. Lastly, ATP leaves through the ATP channel and out of the mitochondria.

Glycolysis is a metabolic pathway that takes place in the cytosol of cells in all living organisms. This pathway can function with or without the presence of oxygen. In humans, aerobic conditions produce pyruvate and anaerobic conditions produce lactate. In aerobic conditions, the process converts one molecule of glucose into two molecules of pyruvate (pyruvic acid), generating energy in the form of two net molecules of ATP. Four molecules of ATP per glucose are actually produced, however, two are consumed as part of the preparatory phase. The initial phosphorylation of glucose is required to increase the reactivity (decrease its stability) in order for the molecule to be cleaved into two pyruvate molecules by the enzyme aldolase. During the pay-off phase of glycolysis, four phosphate groups are transferred to ADP by substrate-level phosphorylation to make four ATP, and two NADH are produced when the pyruvate are oxidized. The overall reaction can be expressed this way:



Starting with glucose, 1 ATP is used to donate a phosphate to glucose to produce glucose 6-phosphate. Glycogen can be converted into glucose 6-phosphate as well with the help of glycogen phosphorylase. During energy metabolism, glucose 6-phosphate becomes fructose 6-phosphate. An additional ATP is used to phosphorylate fructose 6-phosphate into fructose 1,6-disphosphate by the help of phosphofructokinase. Fructose 1,6-diphosphate then splits into two phosphorylated molecules with three carbon chains which later degrades into pyruvate.

Glycolysis can be literally translated as "sugar splitting".^[3]

Oxidative decarboxylation of pyruvate

Main article: Pyruvate decarboxylation

Pyruvate is oxidized to acetyl-CoA and CO₂ by the pyruvate dehydrogenase complex (PDC). The PDC contains multiple copies of three enzymes and is located in the mitochondria of eukaryotic cells and in the cytosol of prokaryotes. In the conversion of pyruvate to acetyl-CoA, one molecule of NADH and one molecule of CO₂ is formed.

Citric acid cycle

Main article: Citric acid cycle

This is also called the *Krebs cycle* or the *tricarboxylic acid cycle*. When oxygen is present, acetyl-CoA is produced from the pyruvate molecules created from glycolysis. Once acetyl-CoA is formed, aerobic or anaerobic respiration can occur.^[4] When oxygen is present, the mitochondria will undergo aerobic respiration which leads to the Krebs cycle. However, if oxygen is not present, fermentation of the pyruvate molecule will occur. In the presence of oxygen, when acetyl-CoA is produced, the molecule then enters the citric acid cycle (Krebs cycle) inside the mitochondrial matrix, and is oxidized to CO₂ while at the same time reducing NAD to NADH. NADH can be used by the electron transport chain to create further ATP as part of oxidative phosphorylation. To fully oxidize the equivalent of one glucose molecule, two acetyl-CoA must be metabolized by the Krebs cycle. Two waste products, H₂O and CO₂, are created during this cycle.

The citric acid cycle is an 8-step process involving 18 different enzymes and co-enzymes.^[4] During the cycle, acetyl-CoA (2 carbons) + oxaloacetate (4 carbons) yields citrate (6 carbons), which is rearranged to a more reactive form called isocitrate (6 carbons). Isocitrate is modified to become α -ketoglutarate (5 carbons), succinyl-CoA, succinate, fumarate, malate, and, finally, oxaloacetate.

The net gain of high-energy compounds from one cycle is 3 NADH, 1 FADH₂, and 1 GTP; the GTP may subsequently be used to produce ATP. Thus, the total yield from 1 glucose molecule (2 pyruvate molecules) is 6 NADH, 2 FADH₂, and 2 ATP.

Oxidative phosphorylation

Main articles: Oxidative phosphorylation, Electron transport chain, Electrochemical gradient, and ATP synthase

In eukaryotes, oxidative phosphorylation occurs in the mitochondrial cristae. It comprises the electron transport chain that establishes a proton gradient (chemiosmotic potential) across the boundary of inner membrane by oxidizing the NADH produced from the Krebs cycle. ATP is synthesized by the ATP synthase enzyme when the chemiosmotic gradient is used to drive the phosphorylation of ADP. The electrons are finally transferred to exogenous oxygen and, with the addition of two protons, water is formed.

Efficiency of ATP production


The table below describes the reactions involved when one glucose molecule is fully oxidized into carbon dioxide. It is assumed that all the reduced coenzymes are oxidized by the electron transport chain and used for oxidative phosphorylation.

Although there is a theoretical yield of 38 ATP molecules per glucose during cellular respiration, such conditions are generally not realized because of losses such as the cost of moving pyruvate (from glycolysis), phosphate, and ADP (substrates for ATP synthesis) into the mitochondria. All are actively transported using carriers that utilize the stored energy in the proton electrochemical gradient.

- Pyruvate is taken up by a specific, low K_m transporter to bring it into the mitochondrial matrix for oxidation by the pyruvate dehydrogenase complex.
- The **phosphate carrier** (PiC) mediates the electroneutral exchange (antiport) of phosphate (H_2PO_4^- ; P_i) for OH^- or symport of phosphate and protons (H^+) across the inner membrane, and the driving force for moving phosphate ions into the mitochondria is the proton motive force.

- The **ATP-ADP translocase** (also called adenine nucleotide translocase, ANT) is an antiporter and exchanges ADP and ATP across the inner membrane. The driving force is due to the ATP (-4) having a more negative charge than the ADP (-3), and thus it dissipates some of the electrical component of the proton electrochemical gradient.

The outcome of these transport processes using the proton electrochemical gradient is that more than 3 H⁺ are needed to make 1 ATP. Obviously this reduces the theoretical efficiency of the whole process and the likely maximum is closer to 28–30 ATP molecules.^[2] In practice the efficiency may be even lower because the inner membrane of the mitochondria is slightly leaky to protons.^[5] Other factors may also dissipate the proton gradient creating an apparently leaky mitochondria. An uncoupling protein known as thermogenin is expressed in some cell types and is a channel that can transport protons. When this protein is active in the inner membrane it short circuits the coupling between the electron transport chain and ATP synthesis. The potential energy from the proton gradient is not used to make ATP but generates heat. This is particularly important in brown fat thermogenesis of newborn and hibernating mammals.



Stoichiometry of aerobic respiration and most known fermentation types in eucaryotic cell.^[6] Numbers in circles indicate counts of carbon atoms in molecules, C₆ is glucose C₆H₁₂O₆, C₁ carbon dioxide CO₂. Mitochondrial outer membrane is omitted.

According to some of newer sources the ATP yield during aerobic respiration is not 36–38, but only about 30–32 ATP molecules / 1 molecule of glucose ^[6], because:

- ATP : NADH+H⁺ and ATP : FADH₂ ratios during the oxidative phosphorylation appear to be not 3 and 2, but 2.5 and 1.5 respectively. Unlike in the substrate-level phosphorylation, the stoichiometry here is difficult to establish.
- ATP synthase produces 1 ATP / 3 H⁺. However the exchange of matrix ATP for cytosolic ADP and Pi (antiport with OH⁻ or symport with H⁺) mediated by ATP–ADP translocase and phosphate carrier consumes 1 H⁺ / 1 ATP as a result of regeneration of the transmembrane potential changed during this transfer, so the net ratio is 1 ATP : 4 H⁺.
- The mitochondrial electron transport chain proton pump transfers across the inner membrane 10 H⁺ / 1 NADH+H⁺ (4 + 2 + 4) or 6 H⁺ / 1 FADH₂ (2 + 4).

So the final stoichiometry is

$$1 \text{ NADH+H}^+ : 10 \text{ H}^+ : 10/4 \text{ ATP} = 1 \text{ NADH+H}^+ : 2.5 \text{ ATP}$$

$$1 \text{ FADH}_2 : 6 \text{ H}^+ : 6/4 \text{ ATP} = 1 \text{ FADH}_2 : 1.5 \text{ ATP}$$

- ATP : NADH+H⁺ coming from glycolysis ratio during the oxidative phosphorylation is
- 1.5, as for FADH₂, if hydrogen atoms (2H⁺+2e⁻) are transferred from cytosolic NADH+H⁺ to mitochondrial FAD by the glycerol phosphate shuttle located in the inner mitochondrial membrane.
- 2.5 in case of malate-aspartate shuttle transferring hydrogen atoms from cytosolic NADH+H⁺ to mitochondrial NAD⁺

So finally we have, per molecule of glucose

- Substrate-level phosphorylation: 2 ATP from glycolysis + 2 ATP (directly GTP) from Krebs cycle

- Oxidative phosphorylation
- 2 NADH+H⁺ from glycolysis: 2 × 1.5 ATP (if glycerol phosphate shuttle transfers hydrogen atoms) or 2 × 2.5 ATP (malate-aspartate shuttle)
- 2 NADH+H⁺ from the oxidative decarboxylation of pyruvate and 6 from Krebs cycle: 8 × 2.5 ATP
- 2 FADH₂ from the Krebs cycle: 2 × 1.5 ATP

Altogether this gives 4 + 3 (or 5) + 20 + 3 = 30 (or 32) ATP per molecule of glucose

The total ATP yield in ethanol or lactic acid fermentation is only 2 molecules coming from glycolysis, because pyruvate is not transferred to the mitochondrion and finally oxidized to the carbon dioxide (CO₂), but reduced to ethanol or lactic acid in the cytoplasm.^[6]

Fermentation

Main article: Fermentation

Without oxygen, pyruvate (pyruvic acid) is not metabolized by cellular respiration but undergoes a process of fermentation. The pyruvate is not transported into the mitochondrion, but remains in the cytoplasm, where it is converted to waste products that may be removed from the cell. This serves the purpose of oxidizing the electron carriers so that they can perform glycolysis again and removing the excess pyruvate. Fermentation oxidizes NADH to NAD⁺ so it can be re-used in glycolysis. In the absence of oxygen, fermentation prevents the buildup of NADH in the cytoplasm and provides NAD⁺ for glycolysis. This waste product varies depending on the organism. In skeletal muscles, the waste product is lactic acid. This type of fermentation is called lactic acid fermentation. In strenuous exercise, when energy demands exceed energy supply, the respiratory chain cannot process all of the hydrogen atoms joined by NADH. During anaerobic glycolysis, NAD⁺ regenerates when pairs of hydrogen

combine with pyruvate to form lactate. Lactate formation is catalyzed by lactate dehydrogenase in a reversible reaction. Lactate can also be used as an indirect precursor for liver glycogen. During recovery, when oxygen becomes available, NAD^+ attaches to hydrogen from lactate to form ATP. In yeast, the waste products are ethanol and carbon dioxide. This type of fermentation is known as alcoholic or ethanol fermentation. The ATP generated in this process is made by substrate-level phosphorylation, which does not require oxygen.

Fermentation is less efficient at using the energy from glucose: only 2 ATP are produced per glucose, compared to the 38 ATP per glucose nominally produced by aerobic respiration. This is because the waste products of fermentation still contain chemical potential energy that can be released by oxidation. Ethanol, for example, can be burned in an internal combustion engine like gasoline. Glycolytic ATP, however, is created more quickly. For prokaryotes to continue a rapid growth rate when they are shifted from an aerobic environment to an anaerobic environment, they must increase the rate of the glycolytic reactions. For multicellular organisms, during short bursts of strenuous activity, muscle cells use fermentation to supplement the ATP production from the slower aerobic respiration, so fermentation may be used by a cell even before the oxygen levels are depleted, as is the case in sports that do not require athletes to pace themselves, such as sprinting.

Anaerobic respiration

Main article: Anaerobic respiration

Cellular respiration is the process by which biological fuels are oxidised in the presence of an inorganic electron acceptor (such as oxygen) to produce large amounts of energy, to drive the bulk production of ATP.

Anaerobic respiration is used by some microorganisms in which neither oxygen (aerobic respiration) nor pyruvate derivatives (fermentation) is the final electron acceptor. Rather, an inorganic acceptor such as sulfate or nitrate is used. Such organisms are typically found in unusual places such as underwater caves or near hydrothermal vents at the bottom of the ocean.

Flavin adenine dinucleotide, or **FADH₂**, is a redox cofactor that is created during the Krebs cycle and utilized during the last part of respiration, the electron transport chain. **Nicotinamide adenine dinucleotide**, or **NADH**, is a similar compound used more actively in the electron transport chain as well. In fact, more NADH is produced and used than FADH₂ in the process of creating energy. There are actually six NADH produced and only two FADH₂ molecules.